

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/341625876>

# The Management of Patients Diagnosed with Incidental Prostate Cancer: Narrative Review

Article in *Research and Reports in Urology* · May 2020

DOI: 10.2147/RRU.S245669

CITATIONS

0

READS

17

6 authors, including:



**Amirreza Abedi**

Shahid Beheshti University of Medical Sciences

33 PUBLICATIONS 92 CITATIONS

SEE PROFILE



**Saleh Ghiasy**

Shahid Beheshti University of Medical Sciences

25 PUBLICATIONS 90 CITATIONS

SEE PROFILE



**Morteza Fallah Karkan**

Shahid Beheshti University of Medical Sciences

78 PUBLICATIONS 282 CITATIONS

SEE PROFILE



**Amirhossein Rahavian**

Shahid Beheshti University of Medical Sciences

20 PUBLICATIONS 32 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Infertility Management [View project](#)



Sexual dysfunction [View project](#)

# The Management of Patients Diagnosed with Incidental Prostate Cancer: Narrative Review

This article was published in the following Dove Press journal:  
*Research and Reports in Urology*

Amir Reza Abedi<sup>1</sup>  
Saleh Ghiasy<sup>1</sup>  
Morteza Fallah-karkan<sup>1,2</sup>  
Amirhossein Rahavian<sup>1,3</sup>  
Farzad Allameh<sup>1,2</sup>

<sup>1</sup>Urology Department, Shohada-e-Tajrish Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran; <sup>2</sup>Center of Excellence for Training Laser Applications in Medicine, Shohada-e-Tajrish Hospital, Shahid Beheshti University of Medical Sciences, Ministry of Health, Tehran, Iran; <sup>3</sup>Infertility and Reproductive Health Research Center (irhrc), Shahid Beheshti Medical Science University, Tehran, Iran

**Abstract:** 5–14% of patients underwent surgery for benign prostate hyperplasia harboring prostate cancer (PCa) focus. The best management of incidental prostate cancer (iPCa) has been debated. The decision “treatment or no treatment” should be determined by predictors which accurately foretell PCa progression after transurethral resection of the prostate (TURP). The purpose of this study is to review the available data that can be useful in daily clinical judgment. Transrectal ultrasound prostate biopsy (TRUSBx) did not provide further Gleason score (GS) data in most patients diagnosed with iPCa. TRUSBX may be useful before active surveillance, but not in all following radical prostatectomy. The decision “treatment or no treatment” should be dependent on the expected chance of having residual cancer and clinical progression. Prostate-specific antigen (PSA) levels before and after TURP are good predictors of residual cancer after TURP. Pathological report of T0 is most likely seen in patients with low PSA density after TURP and indistinguishable lesion on multiparametric magnetic resonance imaging. The decision “treatment vs no treatment” is judged by life expectancy, tumor characteristic in the pathology report of TURP sample and PSA level following TURP. Active surveillance should be contemplated in patients with iPCa who have both prostate-specific antigen density  $\leq 0.08$  after TURP and indistinguishable cancer lesion on multiparametric magnetic resonance imaging. Patients who do not meet the criteria for active surveillance are candidates for radical prostatectomy or radiotherapy (RT). Radical prostatectomy could be peacefully done after TURP with somewhat greater morbidity. RT in patients who had a history of TURP could be safely done and is associated with acceptable quality of life.

**Keywords:** incidental prostate cancer, prostate cancer, transurethral resection of the prostate, TURP

## Introduction

Transurethral resection of the prostate (TURP) is the best option for benign prostatic hyperplasia surgery.<sup>1</sup> The rate of iPCa between patients undergoing TURP without prior diagnosis is between 5% and 14%.<sup>2–5</sup>

According to the percentage of cancerous tissue resected after TURP, iPCa can be subclassified as clinical stage T1a or T1b.<sup>6</sup> Although most of the incidental prostate cancers are found clinically insignificant, recent studies have proposed that in some of them, the clinical course becomes aggressive.<sup>7,8</sup>

The best management of iPCa has been questioned for decades.<sup>9,10</sup> Active surveillance (AS) for every patient with incidental T1 PCa after TURP is not acceptable. The decision “treatment or no treatment” should be chosen by predictors that accurately predict PCa progression after TURP.<sup>11</sup> Radical prostatectomy (RP) after previous

Correspondence: Farzad Allameh  
Department of Urology, Shohada-e-Tajrish Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran  
Tel/Fax +98 2122736386  
Email farzadallame@gmail.com

TURP can be technically challenging.<sup>12,13</sup> It is difficult to recognize those patients at high risk of cancer progression and subsequently need treatment.

In this study, we review other studies to figure out how to manage patients with iPCa after TURP.

## Method

We searched the PubMed database from 1980 to 2019 using the following key words: incidental, prostate cancer, TURP. This was done in order to ensure the comprehensive inclusion of articles related to all incidental prostate cancer. The initial search resulted in 182 articles. Special emphasis was given to relevant articles reporting the management and outcomes of incidental prostate cancer (T1a or T1b). All English papers were included and non-English papers were excluded. According to the predefined inclusion and exclusion criteria, 155 articles were excluded. References from the included studies were manually retrieved to identify additional studies of interest.

## Definition and Epidemiology

iPCa is described as a tumor diagnosed incidentally after surgery for benign prostate hyperplasia (without previous suspicion of PCa) or found after autopsy or detected incidentally after radical cystoprostatectomy for bladder cancer.<sup>14</sup>

Regarding pathology report, incidental PCa after TURP is divided into two groups (clinical T1a or T1b) dependent on the percentage of tumor in the specimen: <5% vs >5%, respectively.<sup>6,15–17</sup> Some study reported cut-off point of 3.8 ng/mL for PSA level to detect iPCa.<sup>18</sup>

## Can We Predict the Possibility of Having Residual Cancer and Clinical Progression?

The decision “treatment or no treatment” should be dependent on the expected chance of having residual cancer and clinical progression.<sup>12</sup> Melchior et al<sup>11</sup> found many patients with incidental carcinoma of prostate after TURP will harbor either no residual cancer or tumors with favorable features in their RP specimens. But, at this time there is no chance to exactly foresee the absence of aggressive PCa after TURP. This dilemma of overtreatment for cancer incidentally detected following TURP versus no treatment with the possibility of tumor progression should be discussed with the patient.

Most urologists recommend transrectal ultrasound-guided needle biopsy (TRUSBx) when iPCa found after

TURP to provide better assessment of residual cancer and tumor grade<sup>2,19</sup> but Lee et al demonstrated that TRUSBx did not provide further Gleason score (GS) data in 86% of patients. They conclude TRUSBx may be useful before AS, but not in all following RP.<sup>19,20</sup>

PSA levels before and after TURP are good predictors of residual cancer after TURP.<sup>21</sup> Umberto Capitation categorized patients into three groups based on the chance of having residual disease at RP and showed that patients with PSA after Surgery for Benign Prostate Hyperplasia (SxBPH) >1.0 ng/mL have the least probability of having T0 but patients with PSA after SxBPH <1.0 ng/mL and PSA before SxBPH <2.0 ng/mL have the highest probability of having residual cancer.<sup>12</sup>

Descazeaud et al<sup>22</sup> demonstrated that patients with iPCa after TURP who have two or more adverse following factors were significantly accompanied with cancer progression: preoperative PSA > or =10 ng/mL, postoperative PSA > or =2 ng/mL, prostate weight > or =60 g, weight of resected tissue > or =40 g, and Gleason score > or =6.

Patients who have low prostate-specific antigen density after TURP and undistinguishable lesion on multiparametric magnetic resonance imaging have the least risk of having residual cancer and should be considered for AS.<sup>23</sup>

PSA before and after TURP and Gleason scores were independent predictors of residual cancer at RP but Stage (T1a vs T1b) but did not predict residual cancer.<sup>12</sup>

Cantrell et al<sup>24</sup> reported that GS is a significant factor to predict clinical progression during follow-up. However, other studies showed that GS is not a good predictor for pathological outcomes after RP or clinical progression.<sup>25,26</sup>

Higher preoperative PSA seemed to be a predictor of iPCa after holmium laser enucleation of the prostate (HoLEP).<sup>27,28</sup> Diabetes may be a key factor to predict the presence of high-risk PCa in men who have undergone HoLEP.<sup>29</sup>

## Treatments

Sebastian et al proposed an individualized approach to counsel the patient who had diagnosed with iPCa after TURP, the decision “treatment vs AS” is judged by patient’s age, life expectancy, tumor characteristic in the pathology report of TURP sample and PSA level following TURP.<sup>11</sup>

1 – AS: should be contemplated for well-differentiated cancers in patients with limited life expectancy and low PSA levels following TURP but a low percentage of patients (up to 21%) may be at hazard for cancer progression.<sup>22</sup>

Patients with iPCa who have both PSA-density ≤0.08 after TURP and indistinguishable cancer lesion on

multiparametric magnetic resonance imaging seem to be considered for AS.<sup>23</sup>

Patients in whom PSA before surgery is less than 2 ng/mL and PSA after surgery is less than 1 ng/mL, the probability of being T0 is high<sup>12</sup> so AS can be proposed to this group.

According to EAU guideline AS or watchful waiting is the best option for patients with T1 PCa if GS is 6 or less and the life expectancy of the patient is less than 10 years.<sup>30</sup> Liu Z et al showed that the clinical result of iPCa was pleasing with the initial treatment of watchful waiting in the Chinese population.<sup>31</sup>

2 – RP: According to EAU guideline RP is the best option for cases with T1b cancer and a life expectancy of more than 10 years.<sup>30</sup> According to AUA guideline AS, brachytherapy, external beam radiotherapy, and RP are suitable options for low-risk PCa.<sup>32</sup>

Colombo et al demonstrated that RP could be peacefully performed after TURP with somewhat greater morbidity.<sup>13</sup>

RP should be recommended in patients with a long life expectancy, poorly differentiated tumors or high PSA levels after TURP.<sup>33,34</sup> RP provides ten-year biochemical-free survival rates as high as 79–98% and 74–88% in T1a and T1b cases, respectively.<sup>12,35,36,37</sup>

Matanhelia et al showed that curative treatments need to be considered for younger cases with a long life expectancy or patients with higher-risk disease.<sup>33</sup>

Paul et al<sup>38</sup> studied 52 cases with iPCa after TURP undergoing RP and showed that TURP is not an adverse predictive factor and morbidity is similar to patients who were detected by needle biopsy. Gacci et al<sup>39</sup> showed that Men who treated with TURP before RP presented an overall incidence of positive surgical margin (PSM) similar to those without previous TURP. However, several studies reported higher PSM rates (21.8–34.2%) in patients who underwent latent-RP (LRP) after TURP.<sup>40,41</sup> Katz et al noted PSM in 12 of 35 patients who underwent LRP after previous TURP<sup>41</sup> Jaffe et al<sup>40</sup> reported a greater overall PSM rate after TURP.

Yang et al<sup>42</sup> showed that LRP is a possible but difficult procedure following TURP. LRP necessitates longer operating times, greater blood loss, higher complication rates and worse short-term continence outcomes.<sup>42</sup> However, controversies occur about the influence of a laparoscopic approach on functional results. Menard et al<sup>43</sup> showed that laparoscopic RP is accompanied with compromised erectile function but the urinary continence rate was not troubled. In contrast, Teber et al<sup>44</sup> reported that the interval to total continence was deferred, with no influence on potency rates. No high level of evidence is

about long-term oncologic and functional results after robot-assisted RP.

Previous TURP distorts the proper surgical plane, which increases the difficulties of later procedures. Elder et al<sup>45</sup> recommended doing surgery either during the first month after TURP or to wait until 4 months after TURP to achieve a satisfactory functional and oncological prognosis for patients with PCa. Zugor et al<sup>46</sup> proposed a time interval between TURP and RP of at least 3 months.

3 – RT: External radiotherapy is an appropriate treatment option in patient with prior history of surgical treatment for BPH and is accompanied with acceptable quality of life. The incidence of severe long-term urinary toxicities is similar to those without a history of TURP.<sup>47</sup> Devisetty et al reported severe GU toxicity in patients who underwent external beam radiotherapy (EBRT) after TURP, but toxicity tends not to persist.<sup>48</sup>

Mélanie Guilhen et al<sup>47</sup> reported External RT is a good option without a major hazard for urinary toxicity in a patient with a history of previous TURP. Lee et al<sup>49</sup> demonstrated that late urinary incontinence occurs in 2% vs 0.2% of patients after RT with or without a history of TURP, respectively. Perez et al<sup>50,51</sup> reported that a non-significant increase in the incontinence rate occurs. Sandhu et al<sup>52</sup> did not find any difference in terms of severe long-term urinary toxicity in patients treated by Three Dimensional (3D) Conformal Radiation Therapy 3DRT or Intensity Modulated Radiotherapy (IMRT) with or without a history of TURP (10 vs 9%). In a systematic review of the literature, four out of 13 studies reported a significantly higher incontinence rate in patients with history of TURP compared to patients without history of TURP. Acute severe urinary toxicity, longer follow-up time, and stage  $\geq$ T3 were shown as adverse parameters for incontinence.<sup>53</sup> Late urinary toxicity rate was diminished in patients treated by modern irradiation techniques such as IMRT.<sup>47</sup> Zapatero et al<sup>54</sup> demonstrated that late urinary complications with high-dose IMRT compared with 3DCRT were lower despite higher radiation dose (80.7 Gy vs 78.7 Gy,  $p < 0.001$ ).

Chevli et al<sup>55</sup> reported that prostate volume does not seem to be a good predictor of RT toxicity in patients with or without TURP. Polland et al<sup>56</sup> showed that older age or pre-TURP urinary urgency seems to be an indicator of post-TURP incontinence in patients who underwent brachytherapy or EBRT for PCa.

## Conclusion

The decision “treatment vs no treatment” is judged by life expectancy, tumor characteristic in the pathology report of

TURP sample and PSA level following TURP. Active surveillance should be contemplated in patients with iPCa who have both prostate-specific antigen density  $\leq 0.08$  after TURP and indistinguishable cancer lesion on multiparametric magnetic resonance imaging. Patients who do not meet the criteria for active surveillance are candidates for radical prostatectomy or radiotherapy (RT). Radical prostatectomy could be peacefully done after TURP with somewhat greater morbidity. RT in patients who had history of TURP could be safely done and is associated with acceptable quality of life.

## Disclosure

The authors report no conflicts of interest in this work.

## References

- Karami H, Hassanzadeh-hadad A, Fallah-karkan M. Comparing monotherapy with tadalafil or tamsulosin and their combination therapy in men with benign prostatic hyperplasia: a randomized clinical trial. *Urol J*. 2016;13(6):2920–2926.
- Jones J, Follis H, Johnson J. Probability of finding T1a and T1b (incidental) prostate cancer during TURP has decreased in the PSA era. *Prostate Cancer Prostatic Dis*. 2009;12(1):57. doi:10.1038/pcan.2008.14
- Zigeuner RE, Lipsky K, Riedler I, et al. Did the rate of incidental prostate cancer change in the era of PSA testing? A retrospective study of 1127 patients. *Urology*. 2003;62(3):451–455. doi:10.1016/S0090-4295(03)00459-X
- Yoo C, Oh CY, Kim SJ, et al. Preoperative clinical factors for diagnosis of incidental prostate cancer in the era of tissue-ablative surgery for benign prostatic hyperplasia: A Korean multi-center review. *Korean J Urol*. 2012;53(6):391–395. doi:10.4111/kju.2012.53.6.391
- Sakamoto H, Matsumoto K, Hayakawa N, et al. Preoperative parameters to predict incidental (T1a and T1b) prostate cancer. *Can Urol Assoc J*. 2014;8(11–12):E815. doi:10.5489/auaj.1890
- Edge S, Byrd D, Compton C, Fritz A, Greene F, Trotti A. *AJCC Cancer Staging Manual*. 7th ed. NY: Springer; 2010.
- Andrén O, Fall K, Franzén L, Andersson S-O, Johansson J-E, Rubin MA. How well does the Gleason score predict prostate cancer death? A 20-year followup of a population based cohort in Sweden. *J Urol*. 2006;175(4):1337–1340. doi:10.1016/S0022-5347(05)00734-2
- Robinson D, Aus G, Bak J, et al. Long-term follow-up of conservatively managed incidental carcinoma of the prostate A multivariate analysis of prognostic factors. *Scand J Urol Nephrol*. 2007;41(2):103–109. doi:10.1080/00365590600991268
- Razzaghi MR, Karkan MF, Ghiasy S, Javanmard B. Laser application in iran urology: a narrative review. *J Lasers Med Sci*. 2018;9(1):1. doi:10.15171/jlms.2018.01
- Ghiasy S, Abedi AR, Moradi A, et al. Is active surveillance an appropriate approach to manage prostate cancer patients with Gleason Score 3+ 3 who met the criteria for active surveillance? *Turkish J Urol*. 2019;45(4):261. doi:10.5152/tud.
- Melchior S, Hadaschik B, Thüroff S, Thomas C, Gillitzer R, Thüroff J. Outcome of radical prostatectomy for incidental carcinoma of the prostate. *BJU Int*. 2009;103(11):1478–1481. doi:10.1111/bju.2009.103.issue-11
- Capitanio U, Scattoni V, Freschi M, et al. Radical prostatectomy for incidental (stage T1a–T1b) prostate cancer: analysis of predictors for residual disease and biochemical recurrence. *Eur Urol*. 2008;54(1):118–125. doi:10.1016/j.eururo.2008.02.018
- Colombo R, Naspro R, Salonia A, et al. Radical prostatectomy after previous prostate surgery: clinical and functional outcomes. *J Urol*. 2006;176(6):2459–2463. doi:10.1016/j.juro.2006.07.140
- Kimura T, Egawa S. Epidemiology of prostate cancer in Asian countries. *Int J Urol*. 2018;25(6):524–531. doi:10.1111/iju.2018.25.issue-6
- Mottet N, Bellmunt J, Briers E, et al. Guidelines on prostate cancer. *Eur Urol*. 2014;65(1):124–137. doi:10.1016/j.eururo.2013.09.046
- Capitanio U, Briganti A, Suardi N, et al. When should we expect no residual tumor (pT0) once we submit incidental T1a-b prostate cancers to radical prostatectomy? *Int J Urol*. 2011;18(2):148–153. doi:10.1111/iju.2011.18.issue-2
- Zhao Z, Ma W, Zeng G, Qi D. PSCA mRNA expression in preoperatively negative prostate biopsies predicts incidental prostate cancer in patients undergoing transurethral resection of the prostate for benign prostatic hyperplasia. *J Surg Oncol*. 2011;104(6):672–678. doi:10.1002/jso.v104.6
- Abedi A-R, Fallah-karkan M, Allameh F, Ranjbar A, Shadmehr A. Incidental prostate cancer: a 10-year review of a tertiary center, Tehran, Iran. *Res Rep Urol*. 2018;10:1. doi:10.2147/RRU.S146159
- Lee LS, Thiruneelakandasivam S, Hong MK, et al. Are transrectal prostate biopsies routinely indicated in patients with incidentally diagnosed prostate cancer following transurethral resection of the prostate for benign disease? *Urol Int*. 2013;91(4):397–403. doi:10.1159/000350898
- Javanmard B, Razzaghi MR, Javanbakht O, Karkan MF, Ghiasy S. Pathological Association Between Radical Prostatectomy and Needle Biopsy Specimen in Prostate Cancer Patients. *Int J Cancer Manag*. 2020;13(1): 1:e91609. doi:10.5812/ijcm.91609
- Visccher D, Cangh V. Assessing the risk of unsuspected prostate cancer in patients with benign prostatic hypertrophy: a 13-year retrospective study of the incidence and natural history of T1a-T1b prostate cancers. *BJU Int*. 1999;84(9):1015–1020. doi:10.1046/j.1464-410x.1999.00386.x
- Descazeaud A, Peyromaure M, Salin A, et al. Predictive factors for progression in patients with clinical stage T1a prostate cancer in the PSA era. *Eur Urol*. 2008;53(2):355–362. doi:10.1016/j.eururo.2007.06.020
- Chung DY, Goh HJ, Koh DH, et al. Clinical significance of multiparametric MRI and PSA density as predictors of residual tumor (pT0) following radical prostatectomy for T1a-T1b (incidental) prostate cancer. *PLoS One*. 2018;13(12):e0210037. doi:10.1371/journal.pone.0210037
- Cantrell BB, DeKlerk DP, Eggleston JC, Boitnott JK, Walsh PC. Pathological factors that influence prognosis in stage A prostatic cancer: the influence of extent versus grade. *J Urol*. 1981;125(4):516–520. doi:10.1016/S0022-5347(17)55092-2
- Epstein JI, Walsh PC, Brendler CB. Radical prostatectomy for impalpable prostate cancer: the Johns Hopkins experience with tumors found on transurethral resection (stages T1A and T1B) and on needle biopsy (stage T1C). *J Urol*. 1994;152(5):1721–1729. doi:10.1016/S0022-5347(17)32370-4
- Epstein JI, Paull G, Eggleston JC, Walsh PC. Prognosis of untreated stage A1 prostatic carcinoma: a study of 94 cases with extended followup. *J Urol*. 1986;136(4):837–839. doi:10.1016/S0022-5347(17)45097-X
- Otsubo S, Yokomizo A, Mochida O, et al. Significance of prostate-specific antigen-related factors in incidental prostate cancer treated by holmium laser enucleation of the prostate. *World J Urol*. 2015;33(3):329–333. doi:10.1007/s00345-014-1310-9
- Bhojani N, Boris RS, Monn MF, Mandeville JA, Lingeman JE. Coexisting prostate cancer found at the time of holmium laser enucleation of the prostate for benign prostatic hyperplasia: predicting its presence and grade in analyzed tissue. *J Endourol*. 2015;29(1):41–46. doi:10.1089/end.2014.0359
- Ohwaki K, Endo F, Shimbo M, Fujisaki A, Hattori K. Comorbidities as predictors of incidental prostate cancer after Holmium laser enucleation of the prostate: diabetes and high-risk cancer. *Aging Male*. 2017;20(4):257–260. doi:10.1080/13685538.2017.1301417

30. Heidenreich A, Aus G, Bolla M, et al. EAU guidelines on prostate cancer. *Eur Urol*. 2008;53(1):68–80. doi:10.1016/j.eururo.2007.09.002
31. Liu Z, Luo C, Hu S, et al. Diagnosis and treatment of T1a-T1b prostate cancer. *J Peking Univ Health Sci*. 2016;48(5):812–816.
32. Sanda MG, Cadeddu JA, Kirkby E, et al. Clinically localized prostate cancer: AUA/ASTRO/SUO guideline. Part I: risk stratification, shared decision making, and care options. *J Urol*. 2018;199(3):683–690. doi:10.1016/j.juro.2017.11.095
33. Heidenreich A, Bellmunt J, Bolla M, et al. EAU guidelines on prostate cancer. Part 1: screening, diagnosis, and treatment of clinically localised disease. *Eur Urol*. 2011;59(1):61–71. doi:10.1016/j.eururo.2010.10.039
34. Karkan MF, Razzaghi MR, Javanmard B, Tayyebiazar A, Ghiyasi S, Montazeri S. Holmium: YAG laser incision of bladder neck contraction following radical retropubic prostatectomy. *Nephro-Urol Mon*. 2020;11(1):e88677. doi:10.5812/numonthly.88677
35. Helfand BT, Mongiu AK, Kan D, et al. Outcomes of radical prostatectomy for patients with clinical stage T1a and T1b disease. *BJU Int*. 2009;104(3):304–309. doi:10.1111/j.1464-410X.2009.08421.x
36. Allameh F, Azghandi S, Karkan MF. Is chemotherapy related with erectile dysfunction in non-urologic cancer patients? *Int J Cancer Manage*. 2018. doi:10.5812/ijcm.82529
37. Allameh F, Rahavian AH, Ghiyasi S. Prevalence of castration success rate in Iranian metastatic prostate cancer patients: A referral center statistics. *Int J Cancer Manag*. 2018 Oct;11(10):e83613. doi:10.5812/ijcm.83613
38. Paul R, Knebel C, Kübler H, Alschibaja M, Günther M, Hartung R. Incidental carcinoma of the prostate: can we and should we recommend radical prostatectomy? *Der Urologe Ausg A*. 2005;44(9):1052,4–8.
39. Gacci M, Simonato A, Lanciotti M, et al. The impact of prior TURP on radical prostatectomy surgical margins: a multicenter analysis. *Urol Int*. 2013;91(1):62–68. doi:10.1159/000346748
40. Jaffe J, Stakhovsky O, Cathelineau X, Barret E, Vallancien G, Rozet F. Surgical outcomes for men undergoing laparoscopic radical prostatectomy after transurethral resection of the prostate. *J Urol*. 2007;178(2):483–487. doi:10.1016/j.juro.2007.03.114
41. Katz R, Borkowski T, Hoznek A, Salomon L, Gettman MT, Abbou CC. Laparoscopic radical prostatectomy in patients following transurethral resection of the prostate. *Urol Int*. 2006;77(3):216–221. doi:10.1159/000094812
42. Yang Y, Luo Y, Hou G-L, et al. Laparoscopic radical prostatectomy after previous transurethral resection of the prostate in clinical T1a and T1b prostate cancer: a matched-pair analysis. *Urol J*. 2015;12(3):2154–2159.
43. Menard J, de la Taille A, Hoznek A, et al. Laparoscopic radical prostatectomy after transurethral resection of the prostate: surgical and functional outcomes. *Urology*. 2008;72(3):593–597. doi:10.1016/j.urology.2008.03.019
44. Teber D, Cresswell J, Ates M, et al. Laparoscopic radical prostatectomy in clinical T1a and T1b prostate cancer: oncologic and functional outcomes—a matched-pair analysis. *Urology*. 2009;73(3):577–581. doi:10.1016/j.urology.2008.09.059
45. Elder JS, Gibbons RP, Correa RJ, Brannen GE. Morbidity of radical perineal prostatectomy following transurethral resection of the prostate. *J Urol*. 1984;132(1):55–57. doi:10.1016/S0022-5347(17)49462-6
46. Zugar V, Labanaris AP, Porres D, Witt JH. Surgical, oncologic, and short-term functional outcomes in patients undergoing robot-assisted prostatectomy after previous transurethral resection of the prostate. *J Endourol*. 2012;26(5):515–519. doi:10.1089/end.2011.0205
47. Guilhen M, Hennequin C, Ouzaid I, et al. Urinary function and quality of life after radiotherapy for prostate cancer in patients with prior history of surgical treatment for benign prostatic hyperplasia. *Radiat Oncol*. 2018;13(1):209. doi:10.1186/s13014-018-1149-0
48. Devisetty K, Zorn KC, Katz MH, Jani AB, Liauw SL. External beam radiation therapy after transurethral resection of the prostate: a report on acute and late genitourinary toxicity. *Int J Radiat Oncol Biol Phys*. 2010;77(4):1060–1065. doi:10.1016/j.ijrobp.2009.06.078
49. Lee WR, Schultheiss TE, Hanlon AL, Hanks GE. Urinary incontinence following external-beam radiotherapy for clinically localized prostate cancer. *Urology*. 1996;48(1):95–99. doi:10.1016/S0090-4295(96)00085-4
50. Perez CA, Lee HK, Georgiou A, Lockett MA. Technical factors affecting morbidity in definitive irradiation for localized carcinoma of the prostate. *Int J Radiat Oncol Biol Phys*. 1994;28(4):811–819. doi:10.1016/0360-3016(94)90100-7
51. Perez CA, Walz BJ, Zivnuska FR, Pilepich M, Prasad K, Bauer W. Irradiation of carcinoma of the prostate localized to the pelvis: analysis of tumor response and prognosis. *Int J Radiat Oncol Biol Phys*. 1980;6(5):555–563. doi:10.1016/0360-3016(80)90382-X
52. Sandhu AS, Zelefsky MJ, Lee HJ, Lombardi D, Fuks Z, Leibel SA. Long-term urinary toxicity after 3-dimensional conformal radiotherapy for prostate cancer in patients with prior history of transurethral resection. *Int J Radiat Oncol Biol Phys*. 2000;48(3):643–647. doi:10.1016/S0360-3016(00)00714-8
53. Ishiyama H, Hirayama T, Jhaveri P, et al. Is there an increase in genitourinary toxicity in patients treated with transurethral resection of the prostate and radiotherapy?: a systematic review. *Am J Clin Oncol*. 2014;37(3):297–304. doi:10.1097/COC.0b013e3182546821
54. Zapatero A, Roch M, Büchser D, et al. Reduced late urinary toxicity with high-dose intensity-modulated radiotherapy using intra-prostate fiducial markers for localized prostate cancer. *Clin Transl Oncol*. 2017;19(9):1161–1167. doi:10.1007/s12094-017-1655-9
55. Chevli C, Narayanan R, Rambarran L, Kubicek G, Chevli KK, Duff M. Effect of pretreatment prostate volume on urinary quality of life following intensity-modulated radiation therapy for localized prostate cancer. *Res Rep Urol*. 2013;5:29.
56. Polland A, Vertosick EA, Sjoberg DD, et al. Preoperative symptoms predict continence after post-radiation transurethral resection of prostate. *Can J Urol*. 2017;24(4):8903–8909.

## Research and Reports in Urology

### Publish your work in this journal

Research and Reports in Urology is an international, peer-reviewed, open access journal publishing original research, reports, editorials, reviews and commentaries on all aspects of adult and pediatric urology in the clinic and laboratory including the following topics: Pathology, pathophysiology of urological disease; Investigation and

treatment of urological disease; Pharmacology of drugs used for the treatment of urological disease. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/research-and-reports-in-urology-journal>

Dovepress